

# An approximate analytical model for footprint estimation of scalar fluxes in thermally stratified atmospheric flows

Cheng-I Hsieh <sup>a</sup>, Gabriel Katul <sup>b,\*</sup>, Tze-wen Chi <sup>c</sup>

<sup>a</sup> *Environment and Ecology, Department of Humanities and General Sciences, National Taiwan University of Science and Technology, Taipei 10660, Taiwan, ROC*

<sup>b</sup> *School of the Environment, Duke University, Box 90328, Durham, NC 27708, USA*

<sup>c</sup> *Department of Environmental Management, Chaoyang University of Technology, Wufeng, Taichung 41342, Taiwan, ROC*

Received 20 May 1999; received in revised form 20 September 1999; accepted 7 October 1999

## Abstract

An approximate analytical model was developed to estimate scalar flux footprint in thermally stratified atmospheric surface layer flows. The proposed model was based on a combination of Lagrangian stochastic dispersion model results and dimensional analysis. The main advantage of this model is its ability to analytically relate atmospheric stability, measurement height, and surface roughness length to flux and footprint. Flux estimation by the proposed model was in good agreement with those calculated by detailed Eulerian and Lagrangian models. Measured water vapor fluxes collected along a downwind transect of a transition from a desert to an irrigated potato site were also used to assess the proposed model performance in the field. It was found that the model well reproduced the measured flux evolution with downwind distance. © 2000 Elsevier Science Ltd. All rights reserved.

## 1. Introduction

Scalar flux footprint estimation continues to be a practical research problem in surface hydrology and boundary layer meteorology [4,11,14,23,27]. Such an analysis is commonly used to quantify the contributing source areas to scalar flux measurement or to examine adequate fetch requirements. Both Eulerian analytical models [6,8] and Lagrangian stochastic dispersion models [5,12,16] with varying degrees of complexity have been used to investigate the relationship between scalar flux and its source areas.

For example (as early as 1986), Gash [6] developed a simple model for footprint calculation for neutral atmospheric conditions. Later, Horst and Weil [7,8] proposed an Eulerian analytical model that is capable of incorporating atmospheric stability, albeit the model treatment of atmospheric stability effects on fetch is not explicit. Leclerc and Thurtell [16] first applied a Lagrangian particle trajectory model to examine “rule of thumb” fetch requirement and found that the “100 to 1 fetch-to-height ratio grossly underestimates fetch re-

quirements when observations are carried out above smooth surfaces, in stable conditions, or at high observation level”. These calculations highlight the important roles of measurement height, surface roughness, and atmospheric stability on footprint estimation. Finn et al. [4] examined the performance of Eulerian [7,8] and Lagrangian [16] models for estimating footprint. They concluded that while Eulerian models are easier to implement, they should be used with caution over rough terrain. In short, despite all such advancements, none of the existing models explicitly describes the relationship between footprint, atmospheric stability, observation height, and surface roughness.

The objective of this study is to develop an approximate analytic expression, analogous to Gash [6], that accurately describes the relationship between footprint, observation height, surface roughness, and atmospheric stability. For this purpose, similarity theory (dimensional analysis) in conjunction with the Lagrangian stochastic dispersion model [25] simulation outputs are used to construct the relationships among parameters (i.e., flux, fetch, atmospheric stability, surface roughness, and observation height) and to develop the model framework. Comparisons with existing Lagrangian and Eulerian models are also performed. The model is also field-tested using water vapor flux measurements

\* Corresponding author. Tel.: +1-919-613-8033; fax: +1-919-684-8741.

E-mail address: gaby@duke.edu (G. Katul).

collected along a progression of distances downwind from the transition of a desert to an irrigated potato field.

## 2. Methods

In this section, a brief review of the Eulerian analytical models proposed by Gash [6] and Horst and Weil [8] and the Lagrangian particle trajectory model of Thomson [25] is provided. A discussion on the performance of the proposed model is then presented.

### 2.1. Gash [6] analytical model

Footprint describes the relative contribution of each upwind surface source, per unit source strength, per unit element, to the measured scalar flux. Mathematically, flux and footprint are related by

$$F(x, z_m) = \int_{-\infty}^x S(x) f(x, z_m) dx, \quad (1)$$

where  $F$  is the scalar flux,  $f$  the footprint,  $S$  ( $\text{g m}^{-2} \text{s}^{-1}$ ) the source strength (or termed as surface flux),  $z_m$  the measurement height, and the mean wind direction is along the horizontal coordinate,  $x$ . Consider a homogeneous velocity field with scalar source strength described as

$$S(x) = \begin{cases} 0 & \text{for } x < 0, \\ S_0 & \text{for } x \geq 0. \end{cases} \quad (2)$$

Gash [6] used Calder's [3] solution for the advection–diffusion equation in a neutral atmosphere and a uniform mean wind field,  $U_u$ , to derive

$$x = -\frac{U_u z_m}{k u_*} \frac{1}{\ln(F/S_0)}, \quad (3)$$

where  $x$  is the fetch requirement (downwind distance) to achieve the desired normalized flux,  $F/S_0$ ,  $u_*$  the friction velocity, and  $k$  ( $= 0.4$ ) is von Karman constant. In (3),  $U_u$  is defined as

$$U_u(z_m) = \frac{\int_{z_0}^{z_m} U(z) dz}{\int_{z_0}^{z_m} dz} = \frac{u_*}{k} (\ln(z_m/z_0) - 1 + z_0/z_m), \quad (4)$$

where  $U(z)$  is the mean wind velocity under neutral condition and is estimated from Monin–Obukov similarity theory (MOST [15,24]), and  $z$  denotes height. By rearranging (3), we have

$$F(x, z_m)/S_0 = \exp\left(-\frac{U_u z_m}{k u_* x}\right) \quad (5)$$

for describing the relationship between flux and fetch. Now, using (1) and (5), we can then estimate footprint by

$$f(x, z_m) = \frac{1}{S_0} \frac{dF(x, z_m)}{dx} = \frac{U_u z_m}{k u_* x^2} \exp\left(-\frac{U_u z_m}{k u_* x}\right) \quad (6)$$

for neutral atmospheric conditions.

### 2.2. Horst and Weil [8] analytical model

With the analytical model for cross-wind-integrated concentration distribution [10,26],

$$C(x, z) = \frac{AQ}{z_p U_p} \exp\left(-\left(z/Bz_p\right)^r\right), \quad (7)$$

Horst and Weil [8] derived

$$f(x, z_m) \approx \left(\frac{dz_p}{dx}\right) \frac{z_m}{z_p^2} \frac{U(z_m)}{U_p(z_p)} A \exp\left(-\left(z_m/Bz_p\right)^r\right) \quad (8)$$

for estimating the footprint, where  $A$  and  $B$  are coefficients defined as  $A = r\Gamma(2/r)/\Gamma^2(1/r)$  and  $B = \Gamma(1/r)/\Gamma(2/r)$ ,  $\Gamma$  the gamma function,  $Q$  is line source strength ( $\text{g m}^{-1} \text{s}^{-1}$ ),  $z_p$  the mean plume height for dispersion from a surface source,  $U_p$  the plume advection velocity, and  $r$  is a concentration profile shape factor. Details for implementing (8) are described in Appendix A.

In the Horst and Weil model, there is no analytical solution for the flux; hence, the flux is calculated by solving (1) numerically.

### 2.3. Lagrangian stochastic dispersion model [25]

The Lagrangian stochastic dispersion model is based on the assumption that the evolution of position and velocity of a fluid element is described by a Markov process.

Basic concepts, details, and many references can be found in [21]. For surface layer dispersion calculation, the horizontal turbulent intensity is typically small so that the evolution of the particle position in the  $x$  direction is generally calculated as [16,18,19]

$$x^{n+1} = x^n + U^n \Delta t, \quad (9a)$$

and the evolution in the  $z$  direction is expressed as

$$z^{n+1} = z^n + w^n \Delta t, \quad (9b)$$

where the superscript  $n$  denotes the  $n$ th time step,  $w$  the instantaneous particle velocity along the  $z$  direction, and  $\Delta t$  is the discrete time interval. By the Markov process assumption, the evolution of  $w$  can be expressed by

$$dw = a(z, w, t) dt + b(z, w, t) d\lambda. \quad (10)$$

In (10),  $a$  is the drift coefficient,  $b$  the random acceleration coefficient, and  $d\lambda$  is a Gaussian random variable with zero mean and variance  $dt$ . For stationary and inhomogeneous Gaussian turbulent flows, by the well-mixed criterion [25] and energy conservation, the unique solutions for  $a$  and  $b$  are:

$$a = -\frac{(w - W)}{t_L} + \frac{1}{2} \frac{\partial \sigma_w^2}{\partial z} \left( 1 + \frac{(w - W)^2}{\sigma_w^2} \right) + \frac{1}{2} \frac{\partial \sigma_w^2}{\partial z} \left( \frac{W(w - W)}{\sigma_w^2} \right) + \frac{\partial W}{\partial z} (W + (w - W)), \quad (11)$$

$$b = \left( \frac{2\sigma_w^2}{t_L} \right)^{1/2}, \quad (12)$$

where  $W$ , the mean Eulerian vertical velocity, is zero in the atmospheric surface layer,  $\sigma_w^2$  the Eulerian vertical velocity variance, and  $t_L$  is the Lagrangian decorrelation time scale. By releasing a large number of particles and using (9a)–(12), and upon specifying the Eulerian velocity statistics profiles and Lagrangian time scale (see Appendix B for details), particle trajectories can be computed, and subsequently, the scalar fluxes and footprint. The scalar flux,  $F$ , at the point  $(x, z_m)$  is calculated as

$$F(x, z_m) = \frac{S_0}{N} (n_\uparrow - n_\downarrow), \quad (13)$$

where  $n_\uparrow$  and  $n_\downarrow$  are the number of particles which reach the height  $z_m$  at position  $x$  with upward and downward vertical velocity, respectively, and  $N$  is the total number of particles released. The footprint,  $f$ , is then calculated from

$$f(x, z_m) = \frac{1}{S_0} \frac{dF(z, z_m)}{dx}.$$

### 2.4. Proposed model

From previous model results [8,12,18,22], the fetch  $x$  is a function of  $F$ ,  $z_m$ ,  $z_0$ , and the atmospheric stability parameter ( $z_m/L$ ), where  $L$  is the Obukhov length (see Appendix A for definition). From (3) and (4)  $z_m$  and  $z_0$  can be combined to form a new length scale,  $z_u$ , defined as

$$z_u = z_m (\ln(z_m/z_0) - 1 + z_0/z_m).$$

Hence, we have three characteristic length scales:  $x$ ,  $L$ , and  $z_u$  for the dimensional analysis. With  $L$  as the key variable, we propose the following two dimensionless groups (Pi groups [20,24]) and write

$$\frac{x}{L} = f\left(\frac{z_u}{L}\right). \quad (14)$$

Using Thomson’s [25] Lagrangian model mentioned above, we calculated the 90% flux fetch requirements (i.e., the  $x$  values for reaching  $F/S_0 = 0.9$ ) for a wide range of  $z_m$ ,  $z_0$ , and  $L$  values. The 90% flux fetch requirement is the needed downwind distance,  $x$  (from the transition), for the measured flux,  $F(x, z_m)$ , to present 90% of the surface flux,  $S_0$ . In our calculations,  $z_m$  ranges from 2 to 20 m;  $z_0$  ranges from 0.01 to 0.1 m;  $L$  ranges from  $-0.1$  to 50 m. Fig. 1(a)–(c) shows how  $x/|L|$




Fig. 1. (a) Scatter plot of  $x/|L|$  versus  $z_u/|L|$  for unstable conditions (top panel); (b) same as (a) but for neutral conditions (middle panel); (c) same as (a) but for stable conditions (bottom panel).

varies with  $z_u/|L|$  for unstable, neutral, and stable atmospheric conditions, respectively. From Fig. 1(a)–(c) and in accordance with (3), we can express  $x/|L|$  as

$$\left(\frac{x}{|L|}\right) = \frac{-1}{k^2 \ln(F/S_0)} D (z_u/|L|)^P, \quad (15)$$

where  $D$  and  $P$  are similarity constants. By applying regression analysis to the results in Fig. 1(a)–(c), we found:

$$\begin{aligned} D = 0.28; P = 0.59 & \quad \text{for unstable condition,} \\ D = 0.97; P = 1 & \quad \text{for near neutral and neutral} \\ & \quad \text{conditions,} \\ D = 2.44; P = 1.33 & \quad \text{for stable condition.} \end{aligned}$$

In [13], near neutral conditions was specified for  $|z/L| < 0.04$ . Here we used a more restricted criterion for near neutral condition with  $|z_u/L| < 0.04$  ( $\approx |z/L| < 0.02$ ). (Note: The relationship between  $z_u/L$  and  $z/L$  is not linear.) Rearranging (15), the flux can be estimated by

$$F(x, z_m)/S_0 = \exp\left(\frac{-1}{k^2 x} D z_u^P |L|^{1-P}\right) \quad (16)$$

and the footprint by

$$f(x, z_m) = \frac{1}{k^2 x^2} D z_u^P |L|^{1-P} \exp\left(\frac{-1}{k^2 x} D z_u^P |L|^{1-P}\right). \quad (17)$$

For neutral conditions, the proposed model reduces to Gash's [6] analytical model with  $D = 0.97$ , which is very close to Gash's analytical value, unity. Upon differentiating  $F(x, z_m)/S_0$  with respect to  $L$ , we obtain

$$\frac{1}{S_0} \frac{dF}{d|L|} = \frac{-1}{k^2 x} (1 - P) D z_u^P |L|^{-P} \exp\left(\frac{-1}{k^2 x} D z_u^P |L|^{1-P}\right), \tag{18}$$

which describes the flux sensitivity to a unit change in the Obukhov length. Differentiating (17) with respect to  $x$  and setting the resultant equation to zero, we obtain

$$x = \frac{D z_u^P |L|^{1-P}}{2k^2}, \tag{19}$$

which permits explicit estimation of the peak location of the footprint as a function of atmospheric stability and  $z_u$ . Eqs. (16)–(19) constitute our proposed model for footprint analysis.

### 3. Model comparisons

The models of Horst and Weil [8], the Lagrangian stochastic dispersion model [25], and our proposed approach for estimating flux and footprint were contrasted for neutral, unstable, and stable atmospheric conditions. For neutral conditions, Gash's [6] model was also considered. A comparison between the proposed model calculation and field measured water vapor flux evolution with downwind distance over a potato site were then discussed.

#### 3.1. Flux and footprint model comparisons

Fig. 2(a) shows a typical comparison among these models for estimating flux as a function of fetch for unstable condition, where  $z_m = 4$  m,  $z_0 = 0.04$  m,  $L = -50$  m. In Fig. 2(a), good agreement is noted among all these models. Fig. 2(b) is the same as Fig. 2(a), but for footprint estimation. Fig. 2(b) shows that the locations of the peak footprint estimated by all three models are reasonably close. Fig. 3(a) and (b) are the same as Fig. 2(a) and (b), respectively, but for neutral condition. All models compared well with each other in both figures. This agreement also demonstrates that Gash's [6] simple model is very reliable in neutral flows when compared to the more detailed models such as those of Horst and Weil [8]. Fig. 4(a) and (b) shows the same comparisons as Fig. 2(a) and (b), respectively, but for stable condition, where  $L = 100$  m. Again, there is good agreement among these models. While footprint peak locations estimated by these models are in close agreement, the magnitudes of the peaks are somewhat different. These differences are neither surprising nor critical since scalar flux magnitude around the peak location changes rapidly within a short distance. In these




Fig. 2. (a) Comparison between flux estimated by the models of Horst and Weil [8] (closed circles), Thomson [25] Lagrangian stochastic (open squares), and the proposed (open circles) as a function of fetch under unstable condition, where  $z_m = 4$  m,  $z_0 = 0.04$  m,  $L = -50$  m (top panel); (b) same as (a) but for footprint (bottom panel).




Fig. 3. (a) Same as Fig. 2(a) but for neutral condition. Prediction by Gash's [6] model (plus) is also shown (top panel); (b) same as Fig. 2(b) but for neutral condition. Prediction by Gash's [6] model (plus) is also shown (bottom panel).




Fig. 4. (a) Same as Fig. 2(a) but for stable condition, where  $L = 100$  m (top panel); (b) same as Fig. 2(b) but for stable condition, where  $L = 100$  m (bottom panel).

calculations, the height-to-fetch ratios are about 1:100, 1:250, and 1:300 for unstable, neutral, and stable conditions, respectively, as shown in Figs. 2(a), 3(a) and 4(a). These results are consistent with those of Leclerc and Thurtell [16].

Of interest is the fetch-to-height ratio, commonly determined from the 90% constant flux layer, where the flux measurements vary within 10% with height. By setting the value of  $F/S_0$  in (15) to be 0.9 and rearranging the equation, we can express

$$\frac{x}{z_m} = \frac{D}{0.105k^2 z_m^{-1} |L|^{1-P} z_u^P} \quad (20)$$

to calculate such fetch-to-height ratio analytically. From (20), it is obvious that the fetch-to-height ratio changes with measurement height, surface roughness, and stability.

Using (18), Fig. 5 shows how flux changes with a unit change in  $L$  at different fetches ( $x$ ) as a function of atmospheric stability, where  $z_m = 4.0$  m and  $z_0 = 0.04$  m. Notice that if the measurement is carried out far away from the leading edge, then the measured flux changes less with stability changes.

Of practical importance is the estimation of contributing source area to a specified measurement level. Using (19), Fig. 6 shows the peak location of the footprint for different measurement heights as a function of atmospheric stability, where  $z_0 = 0.04$  m. It is obvious that




Fig. 5. Flux change,  $(1/S_0)(dF/d|L|)$ , at different fetches ( $x$ ) as a function of atmospheric stability, where  $z_m = 4.0$  m and  $z_0 = 0.04$  m for  $x = 100$  m (plus),  $x = 500$  m (open circles), and  $x = 1000$  m (open squares).

for very unstable conditions, the more co-located the peak location is to the measurement point.

### 3.2. Field testing

Field testing of these footprint models by single point measurements has been conducted by many investigators [4,12]. Here we test the proposed model with field measurements along a progression of distances downwind a transition from a desert to a potato field. A brief description of this experiment is presented below; details can be found in [1].

The experimental site was an irrigated potato field, which was surrounded by a desert and an *Alfalfa* patch. The crop was irrigated frequently to make sure that the crop did not suffer any water deficit. The surface roughness height was 0.005 m for the desert and 0.05 m for the potato field. A stationary eddy-covariance system set at 800 m downwind from the transition was used to measure turbulent fluxes of momentum, sensible heat, latent heat, and  $CO_2$  above the potato field. A mobile eddy-covariance system was used to measure these fluxes along a progression of distances, 1, 38.4, 72, 91, 136, and 295 m, downwind from the transition. Both systems




Fig. 6. Peak location of the footprint at different measurement heights ( $z$ ) as a function of stability, where  $z_0 = 0.04$  m for  $z = 2$  m (plus),  $z = 4$  m (open circles),  $z = 5$  m (open squares),  $z = 8$  m (diamonds), and  $z = 10$  m (stars).

were set at 4 m above the ground and all the data were normalized by the measurements from the stationary system. Field measurements and predictions from a second-order closure model [1] for scalar transport showed that the air was in equilibrium with the potato site at the position of the stationary system. Field measurements also showed that the velocity statistics equilibrated with the potato site over a very short distance from the leading edge. Hence, in a first-order analysis of scalar transport, the velocity statistics are assumed to be planar homogeneous.

At this site, the surface flux upwind is not zero and (2) is not directly applicable. Here the source strength is simply approximated as

$$S(x) = \begin{cases} S_1 & \text{for } x < 0, \\ S_2 & \text{for } x \geq 0, \end{cases} \quad (21)$$

where the leading edge is at  $x = 0$  and  $S_1$  and  $S_2$  are determined from the measured fluxes at  $x = 1$  m and  $x = 800$  m, respectively. By superposition, we can calculate the flux using

$$F(x, z) = S_1 \int_{-\infty}^{-x} f(x, z) dx + S_2 \int_0^x f(x, z) dx. \quad (22)$$




Fig. 7. Variation of the water vapor flux with fetch (downwind distance) above the potato site; the solid line denotes proposed model predictions; the dots denote eddy-correlation measurements.

In (22), for  $x \rightarrow 0$ ,  $F(x, z) \rightarrow S_1$ ; for  $x \rightarrow \infty$ ,  $F(x, z) \rightarrow S_2$ . Using (17), (21) and (22), Fig. 7 shows the variation of water vapor flux with fetch (downwind distance) above the potato site; the solid line denotes the proposed model predictions; the dots denote ensemble-averaged eddy-covariance measurements. In Fig. 7, the predicted water vapor flux variation compared reasonably well with the observation. In this field experiment, the air water vapor concentration increases with distance downwind the transition; thus, the surface water vapor flux will decrease with distance from the interface. Hence, the real source strengths (surface fluxes) along the downwind distance from the discontinuity are stronger than the simple approximation in (21). This perhaps explains why the observed fluxes increase more rapidly downwind of the interface than the model prediction.

#### 4. Conclusion

An approximated analytical model to estimate footprint as a function of atmospheric stability and the length scale  $z_u$  was proposed and field-tested. This model is based on dimensional analysis and output results from a Lagrangian stochastic dispersion model. The model performance is comparable to detailed Eulerian and Lagrangian models. Also, good agreement between measured and model predicted water vapor flux evolution with downwind distance over a potato site was demonstrated.

#### Acknowledgements

The authors are grateful to Dennis Baldocchi for the field measurements. The authors would like to thank the

two anonymous reviewers for their helpful comments. This project was funded, in part, by the National Science Council, Taiwan, and the US Department of Energy's (DOE) National Institute of Global Environmental Change (NIGEC) through the NIGEC Southeast Regional Center at the University of Alabama, Tuscaloosa, under the cooperative agreement DE-FC03-90-ER61010 and the FACE-FACTS Project (DE-FG05-95ER62083).

**Appendix A. The model of Horst and Weil [8]**

To implement (8) for footprint analysis,  $U_p(z_p)$  and the relation between  $x$  and  $z_p$  are needed. With  $z_p$  calculated by the Lagrangian similarity equation [26]

$$\frac{dz_p}{dx} = \frac{k^2}{(\ln(pz_p/z_0) - \psi_m(pz_p/L)) \phi_h(pz_p/L)} \tag{A.1}$$

and Businger–Dyer [2,15] formulas for the stability correction function for velocity ( $\psi_m$ ) and heat ( $\phi_h$ ) profiles, Horst and Weil [8] derived

$$\frac{x}{z_0} = \Psi(z_p) - \Psi(z_0). \tag{A.2}$$

In (A1),  $L$  is the Obukhov length defined as

$$L = \frac{-u_*^3 T}{kg \langle w\theta \rangle}, \tag{A.3}$$

where  $g$  ( $= 9.8 \text{ m s}^{-2}$ ) is the gravitational constant,  $T$  the mean air temperature (in K),  $\theta$  the temperature fluctuation, and  $\langle \rangle$  denotes averaging. The functions in (A.1) and (A.2) are defined as:

$$\phi_h = (1 + \beta z/L), \tag{A.4}$$

$$\Psi_m = (-\beta z/L), \tag{A.5}$$

$$\Psi(z) = \frac{1}{k^2} \frac{z}{z_0} \left\{ \ln(pz/z_0) - 1 + \beta pz/L(1/4 + \beta pz/(3L)) + (1/2) \ln(pz/z_0) \right\} \tag{A.6}$$

for  $z/L \geq 0$  and

$$\phi_h = (1 - \alpha z/L)^{-1/2}, \tag{A.7}$$

$$\Psi_m(y) = 2 \ln((y+1)/2) + \ln((y^2+1)/2) + 2 \tan^{-1}((1-y)/(1+y)), \tag{A.8}$$

$$y = (1 - \alpha z/L)^{1/4}, \tag{A.9}$$

$$\Psi(z) = \frac{1}{k^2} \frac{2|L|}{\alpha pz_0} \left\{ y_p^2 (\ln(pz/z_0) - \psi_m(y_p)) + 2 \tan^{-1}(y_p) + \ln((y_p+1)/(y_p-1)) - 4y_p \right\}, \tag{A.10}$$

$$y_p = (1 - \alpha pz/L)^{1/4} \tag{A.11}$$

for  $z/L < 0$ . Note (A.10) is derived by Horst and Weil [9].

The plume advection velocity,  $U_p$ , is calculated as

$$U_p(Z) = \frac{u_*}{k} (\ln(cz/z_0) - \psi_m(cz/L)). \tag{A.12}$$

The constants in the model of Horst and Weil are:  $\alpha = 16$ ,  $\beta = 5$ ,  $p = 1.55$ , and  $c = 0.66$ ,  $r = 2$  for  $z/L < 0$ ;  $c = 0.63$ ,  $r = 1.5$  for  $z/L = 0$ ;  $c = 0.56$ ,  $r = 1$  for  $z/L > 0$ .

**Appendix B. Lagrangian stochastic dispersion model**

To drive the Lagrangian model, the profiles of Eulerian velocity statistics and Lagrangian time scale need to be specified a priori. By MOST [15], the horizontal mean wind profile,  $U(z)$ , is calculated as

$$U(z) = \frac{u_*}{k} (\ln(z/z_0) - \Psi_m(z/L)), \tag{B.1}$$

where  $\Psi_m$  is taken as (A.5) for  $z/L \geq 0$  and (A.8) for  $z/L < 0$ , and the vertical velocity standard deviation profile,  $\sigma_w$ , is expressed as

$$\sigma_w = 1.25u_*(1 - 3z/L)^{1/3} \text{ for } z/L < 0, \tag{B.2}$$

$$\sigma_w = 1.25u_* \text{ for } z/L \geq 0. \tag{B.3}$$

The Lagrangian time scale,  $t_L$ , is estimated as

$$t_L = \frac{kzu_*}{\phi_h \sigma_w^2} \tag{B.4}$$

to match the dispersion coefficient given by K-theory [17]. The discrete time interval,  $\Delta t$ , is set to be  $0.1t_L$ . Here  $\phi_h$  is estimated by [12]:

$$\phi_h = 0.32(0.037 - z/L)^{-1/3} \text{ for } z/L < 0, \tag{B.5a}$$

$$\phi_h = 1 + 5z/L \text{ for } z/L \geq 0. \tag{B.5b}$$

In the Lagrangian model, surface roughness height,  $z_0$ , is set as a reflection boundary. Tracer particles reaching this boundary are perfectly reflected [28].

**References**

- [1] Baldocchi D, Rao S. Intra-field variability of scalar flux densities across a transition between a desert and an irrigated potato site. *Boundary-Layer Meteor* 1995;76:109–36.
- [2] Businger JA. A note on the Businger–Dyer profiles. *Boundary-Layer Meteor* 1988;42:145–51.
- [3] Calder KL. Some recent British work on the problem of diffusion in the lower atmosphere. In: *Proceedings of the US Tech. Conf. on Air Poll.* New York: McGraw-Hill, 1952. p. 787–92.
- [4] Finn D, Lamb B, Leclerc M, Horst TW. Experimental evaluation of analytical and Lagrangian surface-layer flux footprint models. *Boundary-Layer Meteor* 1996;80:283–308.

- [5] Flesch TK. The footprint for flux measurements, from backward Lagrangian stochastic models. *Boundary-Layer Meteorol* 1996;78:399–404.
- [6] Gash JHC. A note on estimating the effect of a limited fetch on micrometeorological evaporation measurements. *Boundary-Layer Meteorol* 1986;35:409–13.
- [7] Horst TW, Weil JC. Footprint estimation for scalar flux measurements in the atmospheric surface layer. *Boundary-Layer Meteorol* 1992;59:279–96.
- [8] Horst TW, Weil JC. How far is far enough? The fetch requirements for micrometeorological measurement of surface fluxes. *J Atmos Oceanic Technol* 1994;11:1018–25.
- [9] Horst TW, Weil JC. How far is far enough? The fetch requirements for micrometeorological measurement of surface fluxes. *J Atmos Oceanic Technol* 1995;12:447.
- [10] Horst TW. Lagrangian similarity modeling of vertical diffusion from a ground-level source. *J Appl Meteor* 1979;18:733–40.
- [11] Horst TW. The footprint for estimation of atmosphere–surface exchange fluxes by profile techniques. *Boundary-Layer Meteorol* 1999;90:171–88.
- [12] Hsieh C-I, Katul GG, Schieldge J, Sigmon JT, Knoerr KK. The Lagrangian stochastic model for fetch and latent heat flux estimation above uniform and non-uniform terrain. *Water Resour Res* 1997;33:427–38.
- [13] Kader BA, Yaglom AM. Mean fields and fluctuation moments in unstable stratified turbulent boundary layers. *J Fluid Mech* 1990;212:637–62.
- [14] Kaharabata SK, Schuepp PH, Ogunjemiyo S, Shen S, Leclerc MY, Desjardins RL, MacPherson JI. Footprint considerations in BOREAS. *J Geophys Res* 1997;102:29113–24.
- [15] Kaimal JC, Finnigan JJ. *Atmospheric boundary layer flows*. Oxford: Oxford University Press, 1994.
- [16] Leclerc MY, Thurtell GW. Footprint prediction of scalar fluxes using a Markovian analysis. *Boundary-Layer Meteorol* 1990;52:247–58.
- [17] Ley AJ, Thomson DJ. A random walk model of dispersion in the diabatic surface layer. *Quart J Roy Meteor Soc* 1983;109:847–80.
- [18] Luhar AK, Rao KS. Source footprint analysis for scalar fluxes measured in the flows over an inhomogeneous surface. In: Gryning S-E, Millan MM, editors. *Air Pollution Modeling and its Application*. New York: Plenum Press, 1994. p. 315–23.
- [19] Luhar AK, Rao KS. Lagrangian stochastic dispersion model simulations of tracer data in nocturnal flows over complex terrain. *Atmos Environ* 1994;28:3417–31.
- [20] Perry RH, Chilton CH, Kirkpatrick SD, editors. *Perry's chemical engineer's handbook*. 4th ed. New York: McGraw-Hill, 1963, pp. 2-87–90.
- [21] Rodean H. *Stochastic Lagrangian models of turbulent diffusion, meteorological monographs*. American Meteorological Society 1996;26(48):84.
- [22] Schuepp PH, Leclerc MY, Macpherson JI, Desjardins RL. Footprint prediction of scalar fluxes from analytical solutions of the diffusion equation. *Boundary-Layer Meteorol* 1990;50:355–73.
- [23] Stannard DI. A theoretically based determination of Bowen-ratio fetch requirements. *Boundary-Layer Meteorol* 1997;83:375–406.
- [24] Stull RB. *An Introduction to boundary layer meteorology*. MA, USA: Kluwer Academic Publishers, 1988.
- [25] Thomson DJ. Criteria for the selection of stochastic models of particle trajectories in turbulent flows. *J Fluid Mech* 1987;180:529–56.
- [26] Van Ulden AP. Simple estimates for vertical diffusion from sources near the ground. *Atmos Environ* 1978;12:2125–9.
- [27] Wilson JD, Swaters GE. The source area influencing a measurement in the planetary boundary: the footprint and the distribution of contact distance. *Boundary-Layer Meteorol* 1991;55:25–46.
- [28] Wilson JD, Flesch TK. Flow boundaries in random-flight dispersion models: enforcing the well-mixed condition. *J Appl Meteorol* 1993;32:1695–707.